

Collegium Pharmaceutical

H.C. Wainwright 28th Annual Global
Investment Conference

September 16, 2026 | Nasdaq: COLL

*Healthier people.
Stronger communities.*

Forward-Looking Statements

This presentation contains forward-looking statements within the meaning of The Private Securities Litigation Reform Act of 1995. We may, in some cases, use terms such as "predicts," "forecasts," "believes," "potential," "proposed," "continue," "estimates," "anticipates," "expects," "plans," "intends," "may," "could," "might," "should" or other words that convey uncertainty of future events or outcomes to identify these forward-looking statements. Examples of forward-looking statements contained in this press release include, among others, projected financial performance, including expected revenue and adjusted EBITDA; statements related to the anticipated benefits of the acquisition of AZSTARYS, including its impact on Collegium’s ADHD portfolio and commercial strategy; statements related to current and future market opportunities for our products and our assumptions related thereto and other statements that are not historic facts. Such statements are subject to numerous important factors, risks and uncertainties that may cause actual events or results, performance, or achievements to differ materially from the company's current expectations, including risks relating to, among others: our ability to realize the anticipated benefits of the AZSTARYS acquisition, including the possibility that the expected benefits from the acquisition will not be realized or will not be realized within the expected time period; the risk that the businesses will not be integrated successfully; significant transaction costs or the acquisition of unknown liabilities; future opportunities and plans for our products, including uncertainty of the expected financial performance of such products; our ability to commercialize and grow sales of our products; our ability to manage our relationships with licensors; the success of competing products that are or become available; our ability to maintain regulatory approval of our products, and any related restrictions, limitations, and/or warnings in the label of our products; the size of the markets for our products, and our ability to service those markets; our ability to obtain reimbursement and third-party payor contracts for our products; the rate and degree of market acceptance of our products; the costs of commercialization activities, including marketing, sales and distribution; changing market conditions for our products; the outcome of any patent infringement or other litigation that may be brought by or against us; the outcome of any governmental investigation related to our business; our ability to secure adequate supplies of active pharmaceutical ingredient for each of our products and manufacture adequate supplies of commercially saleable inventory; our ability to obtain funding for our operations and business development; regulatory developments in the U.S.; our expectations regarding our ability to obtain and maintain sufficient intellectual property protection for our products; our ability to comply with stringent U.S. and foreign government regulation in the manufacture of pharmaceutical products, including U.S. Drug Enforcement Agency compliance; our customer concentration; and the accuracy of our estimates regarding expenses, revenues, capital requirements and need for additional financing. These and other risks are described under the heading "Risk Factors" in our Annual Reports on Form 10-K and Quarterly Reports on Form 10-Q and other filings with the SEC. Any forward-looking statements that we make in this press release speak only as of the date of this press release. We assume no obligation to update our forward-looking statements whether as a result of new information, future events or otherwise, after the date of this press release.

Non-GAAP Financial Measures

To supplement our financial results presented on a GAAP basis, we have included information about certain non-GAAP financial measures. We believe the presentation of these non-GAAP financial measures, when viewed with our results under GAAP and the accompanying reconciliations, provide analysts, investors, lenders, and other third parties with insights into how we evaluate normal operational activities, including our ability to generate cash from operations, on a comparable year-over-year basis and manage our budgeting and forecasting. In addition, certain non-GAAP financial measures, primarily Adjusted EBITDA, are used to measure performance when determining components of annual compensation for substantially all non-sales force employees, including senior management.

In this presentation, we discuss the following financial measures that are not calculated in accordance with GAAP, to supplement our consolidated financial statements presented on a GAAP basis.

Adjusted EBITDA

Adjusted EBITDA is a non-GAAP financial measure that represents GAAP net income or loss adjusted to exclude interest expense, interest income, the benefit from or provision for income taxes, depreciation, amortization, stock-based compensation, and other adjustments to reflect changes that occur in our business but do not represent ongoing operations. Adjusted EBITDA, as used by us, may be calculated differently from, and therefore may not be comparable to, similarly titled measures used by other companies.

There are several limitations related to the use of adjusted EBITDA rather than net income or loss, which is the nearest GAAP equivalent, such as:

- adjusted EBITDA excludes depreciation and amortization, and, although these are non-cash expenses, the assets being depreciated or amortized may have to be replaced in the future, the cash requirements for which are not reflected in adjusted EBITDA;
- adjusted EBITDA does not reflect changes in, or cash requirements for, working capital needs;
- adjusted EBITDA does not reflect the benefit from or provision for income taxes or the cash requirements to pay taxes;
- adjusted EBITDA does not reflect historical cash expenditures or future requirements for capital expenditures or contractual commitments;
- we exclude stock-based compensation expense from adjusted EBITDA although: (i) it has been, and will continue to be for the foreseeable future, a significant recurring expense for our business and an important part of our compensation strategy; and (ii) if we did not pay out a portion of our compensation in the form of stock-based compensation, the cash salary expense included in operating expenses would be higher, which would affect our cash position;
- we exclude impairment expenses from adjusted EBITDA and, although these are non-cash expenses, the asset(s) being impaired may have to be replaced in the future, the cash requirements for which are not reflected in adjusted EBITDA;
- we exclude restructuring expenses from adjusted EBITDA. Restructuring expenses primarily include employee severance and contract termination costs that are not related to acquisitions. The amount and/or frequency of these restructuring expenses are not part of our underlying business;
- we exclude litigation settlements and contingencies that are subject to recovery from adjusted EBITDA, as well as any applicable income items, credit adjustments, or recoveries due to subsequent changes in estimates. This does not include our legal fees to defend claims, which are expensed as incurred;
- we exclude acquisition-related expenses as these expenses would not have otherwise been incurred in the periods presented as part of our continuing operations and their exclusion is useful to investors because it enhances comparability of operating performance across periods. Acquisition-related expenses include transaction costs, which primarily consisted of financial advisory, banking, legal, and regulatory fees, and other consulting fees, incurred to complete the acquisition, employee-related expenses (severance cost and benefits) for terminated employees after the acquisition, legal defense expenses for specific acquired claims that relate to acts that occurred prior to our acquisition, and miscellaneous other acquisition-related expenses incurred;
- we exclude recognition of the step-up basis in inventory from acquisitions (i.e., the adjustment to record inventory from historic cost to fair value at acquisition) as the adjustment does not reflect the ongoing expense associated with sale of our products as part of our underlying business;
- we exclude losses on extinguishments of debt as these expenses are episodic in nature and do not directly correlate to the cost of operating our business on an ongoing basis;
- we exclude executive transition expenses from adjusted EBITDA as the amount and/or frequency of these expenses are episodic in nature and do not directly correlate to the cost of operating our business on an ongoing basis; and
- we exclude other expenses, from time to time, that are episodic in nature and do not directly correlate to the cost of operating our business on an ongoing basis.

Adjusted Operating Expenses

Adjusted operating expenses is a non-GAAP financial measure that represents GAAP operating expenses adjusted to exclude stock-based compensation expense, and other adjustments to reflect changes that occur in our business but do not represent ongoing operations.

Adjusted Net Income and Adjusted Earnings Per Share

Adjusted net income is a non-GAAP financial measure that represents GAAP net income or loss adjusted to exclude significant income and expense items that are non-cash or not indicative of ongoing operations, including consideration of the tax effect of the adjustments. Adjusted earnings per share is a non-GAAP financial measure that represents adjusted net income per share. Adjusted weighted-average shares - diluted is calculated in accordance with the treasury stock, if-converted, or contingently issuable accounting methods, depending on the nature of the security.

Reconciliations of adjusted EBITDA to the most directly comparable GAAP financial measures are included in this presentation.

The Company has not provided a reconciliation of its full-year 2026 guidance for adjusted EBITDA to the most directly comparable forward-looking GAAP measures, in reliance on the unreasonable efforts exception provided under Item 10(e)(1)(i)(B) of Regulation S-K, because the Company is unable to predict, without unreasonable efforts, the timing and amount of items that would be included in such a reconciliation, including, but not limited to, stock-based compensation expense, acquisition related expenses, and litigation settlements. These items are uncertain and depend on various factors that are outside of the Company’s control or cannot be reasonably predicted. While the Company is unable to address the probable significance of these items, they could have a material impact on GAAP net income and operating expenses for the guidance period. A reconciliation of adjusted EBITDA would imply a degree of precision and certainty as to these future items that does not exist and could be confusing to investors.



Company Overview

Vikram Karnani

President & Chief Executive Officer

Building a Leading, Diversified Biopharmaceutical Company

Healthier people.
Stronger communities.



BY THE NUMBERS

\$840M

2026 Product Sales¹

\$458M

2026 Adjusted
EBITDA^{1,2}

2

Current focus areas:
ADHD & Pain

DIFFERENTIATED MEDICINES

Jornay^{PM}
methylphenidate HCl (II)
extended-release capsules

azstarys[®]
serdexmethylphenidate
and dexmethylphenidate
26.1/5.2mg • 39.2/7.8mg • 52.3/10.4mg capsules

Xtampza[®] ER
(oxycodone) EXTENDED-RELEASE
CAPSULES (II)

BELBUCA[®] (II)

NUCYNTA[®]
(tapentadol) TABLETS (II)

NUCYNTA[®] ER
(tapentadol) EXTENDED-RELEASE
TABLETS (II)

Path to Building a Leading, Diversified Biopharmaceutical Company



History of becoming the leader in responsible pain management



Beginning of diversification strategy



2002

Founded to address **opioid epidemic** with **abuse-deterrent pain medicines**

1990s – 2000s

Rise of opioid epidemic in the U.S.

2016

FDA approved **Xtampza® ER** formulated with proprietary abuse-deterrent technology, **DETERx®**

2015

Initial Public Offering

2018 – 2020

In-licensed and acquired the **Nucynta Franchise**, propelling Collegium into profitability

2022

Acquired **Belbuca®**, solidifying leadership in responsible pain management


2024

Acquired **JORNAY PM®**, expanding commercial presence to neuropsychiatry

2026

Acquired **AZSTARYS®**, accelerating growth trajectory and strengthening position in ADHD

Collegium's Vision for the Next Phase of Growth



2026 Financial Guidance

	2025 Actuals	2026 Guidance Range ²	YoY Change ³
Product Revenues, Net	\$781M	\$825 – 855M	+8%
JORNAY PM Revenue, Net	\$149M	\$190 - 200M	+31%
AZSTARYS Revenue, Net	N/A ⁴	\$65 – 75M	N/A
Adjusted EBITDA ¹	\$460M	\$445 – 470M	Flat

Strategic Priorities to Drive Value Creation



Recent Business Highlights¹

Accelerated Commercial Momentum

ADHD Portfolio



+41% YoY growth
in Q2'26 net revenue



\$12.9 million
in net revenue in partial
quarter of sales to
Collegium

Pain Portfolio



\$140.9 million
in Q2'26 net revenue

Achieved Top-and Bottom-line Growth

Product Revenues, Net

+6%

\$188M

Q2'25

\$200M

Q2'26

Adjusted EBITDA²

+8%

\$105M

Q2'25

\$114M

Q2'26

Strategically Deployed Capital to Expand and Diversify Portfolio




Completed acquisition of AZSTARYS and
successfully integrated into ADHD portfolio

2.1x net debt to adjusted EBITDA
as of June 30, 2026²

**Announced \$50M Accelerated Share Repurchase
(ASR) program** on August 13, 2026

Driving Significant Growth in ADHD Portfolio

Overview of U.S. ADHD Market and Key Commercial Dynamics



Commercial Dynamics

- 1. Large and growing market**
8% CAGR from 2020-2025
- 2. High patient unmet need**
Patients on average try ~3 different ADHD medicines before finding right treatment²
- 3. Stimulants remain the preferred treatment for vast majority of patients**
- 4. Significant opportunity for both JORNAY PM and AZSTARYS to grow share from generic stimulants**
including methylphenidates and amphetamines

Highly Differentiated Medicine in the ADHD Market



- **Highly differentiated** central nervous system (CNS) stimulant prescription medicine for the treatment of ADHD in people six years of age and older in the U.S.
- Only stimulant ADHD medication with **once-daily evening dosing** that provides **symptom control upon awakening**, eliminating need to dose in the morning and wait for onset of action
- **Smooth symptom control throughout the day**, which may eliminate the need for short-acting stimulant add-ons
- Slow absorption in colon, **providing smooth onset and offset of effect**

Product Differentiation and Strong Brand Fundamentals Drive Utilization¹

Jornay^{PM}
methylphenidate HCl 
extended-release capsules

#1 highest rated

branded ADHD medicine in terms of product differentiation

~70%

of surveyed HCPs plan to increase prescribing (highest among all other branded ADHD medicines)

>70%

of HCPs will honor a patient/caregiver request to try JORNAY PM

~67%

unaided recall of JORNAY PM, a significant improvement from Q2'25




#1 highest rated company in terms of reputation in ADHD

Fastest Growing Stimulant for Treatment of ADHD


GROWTH IN QUARTERLY PRESCRIPTIONS¹

+13.1%




STRONG AND GROWING PRESCRIBER BASE²

+17.6%



MARKET SHARE IN BRANDED LONG-ACTING METHYLPHENIDATE MARKET¹

+5.8
Percentage
Points



AZSTARYS is Well Positioned for Future Growth and is a Highly Complementary Addition to Our ADHD Portfolio



- **First** and **only** ADHD treatment with both **fast** and **long-acting** medicines in one capsule
- Treatment option for patients in need of **rapid onset** of efficacy and **duration** throughout the day
- Viewed as **highly differentiated** and **highly favorable** amongst HCPs¹

- **IP protection through 2037**
- **Enables significant cost synergies**
- **Immediately accretive to adjusted EBITDA² with greater impact expected in 2027+**

\$65-75 million in revenue expected between May 12 and December 31, 2026

2026 Priorities: Driving Further Growth for ADHD Portfolio

jornay^{PM}
methylphenidate HCl [Ⓢ]
extended-release capsules

- Increase awareness and adoption with expanded set of prescribers
- Raise caregiver and patient awareness to drive HCP request
- Increase depth of prescribing with targeted physicians

azstar^{ys} [Ⓢ]
serdexmethylphenidate
and dexamethylphenidate
26.1/5.2mg • 39.2/7.8mg • 52.3/10.4mg capsules

- Successfully integrate AZSTARYS into portfolio
- Accelerate growth trajectory by leveraging established commercial infrastructure and expertise
- Evaluate opportunities to drive operational efficiencies

Maintain broad patient access

Maximizing the Durability of the Pain Portfolio

The Leader in Responsible Pain Management

STRONG BRAND FUNDAMENTALS¹


• Belbuca:

- The **ONLY** long-acting opioid pain medicine that uses buprenorphine buccal film technology
- **#1** highest rated branded ER opioid in terms of product differentiation and favorability


• Xtampza:

- The **ONLY** extended-release oxycodone pain medicine that uses best-in-class abuse deterrent technology (DETERx)
- **#1** highest rated ER oxycodone in terms of product differentiation and favorability

PAIN PRODUCT REVENUES, NET²



Strong IP Management of Patent Protected Portfolio



Strategically Deploying Capital and Creating Shareholder Value

Track Record of Successful Business Development Driving Top- and Bottom-Line Growth

Nucynta Franchise (Feb. 2020)

NUCYNTA[®] (tapentadol) TABLETS II **NUCYNTA[®] ER** (tapentadol) EXTENDED-RELEASE TABLETS II

BDSI (Mar. 2022)


biodelivery SCIENCES **BELBUCA[™]** (buprenorphine) Buccal Film

Ironshore Therapeutics (Sep. 2024)

IRONSHORE PHARMACEUTICALS **jornay^{PM}** methylphenidate HCl II extended-release capsules

AZSTARYS (May 2026)

azstarys[®] serdexmethylphenidate and dexamethylphenidate 26.1/5.2mg • 39.2/7.8mg • 52.3/10.4mg capsules



Disciplined Business Development Approach

Guiding Framework for Near-term BD Efforts



TARGET THERAPEUTIC AREAS

- **Neuropsychiatry and pediatrics**
- **Other specialty conditions**
(case-by-case)
- **Rare diseases**
(case-by-case)

ADDITIONAL FEATURES

- Commercial or near-commercial
- Cost efficient sales and marketing requirements
- LOE into 2030's and beyond


While maintaining robust cash generation and financial strength

Opportunistic Share Repurchases Deliver Value to Shareholders¹

*Repurchased 10.7M shares
at average price of \$24.75*

2021 -	2.2M shares at \$19.93
2022 -	1.1M shares at \$17.57
2023 -	3.1M shares at \$24.29
2024 -	1.9M shares at \$31.88
2025 -	0.8M shares at \$30.41
2026 -	1.6M shares at \$25.70²

Returned \$272M to Shareholders since 2021



Summary

Serving Patients and Creating Value for Shareholders

2026 STRATEGIC PRIORITIES

1. Drive further growth for the ADHD Portfolio

2. Maximize the durability of the Pain Portfolio

3. Strategically deploy capital

- Business Development
- Debt repayment
- Share repurchases

VALUE CREATION



Grow
Revenues



Extend
longevity



Increase
profitability



Generate
robust cash flows



Diversify
portfolio



Strengthen
balance sheet

Important Safety Information

Important Safety Information about JORNAY PM (methylphenidate HCl extended-release capsules)

JORNAY PM
(methylphenidate HCl
extended-release
capsules)

WARNING: ABUSE, MISUSE, AND ADDICTION

JORNAY PM has a high potential for abuse and misuse, which can lead to the development of a substance use disorder, including addiction. Misuse and abuse of CNS stimulants, including JORNAY PM, can result in overdose and death, and this risk is increased with higher doses or unapproved methods of administration, such as snorting or injection.

Before prescribing JORNAY PM, assess each patient's risk for abuse, misuse, and addiction. Educate patients and their families about these risks, proper storage of the drug, and proper disposal of any unused drug. Throughout JORNAY PM treatment, reassess each patient's risk of abuse, misuse, and addiction and frequently monitor for signs and symptoms of abuse, misuse and addiction.

CONTRAINDICATIONS

- Known hypersensitivity to methylphenidate or other components of JORNAY PM. Hypersensitivity reactions such as angioedema and anaphylactic reactions have been reported in patients treated with methylphenidate products.
- Concurrent treatment with a monoamine oxidase inhibitor (MAOI), or use of an MAOI within the preceding 14 days because of the risk of hypertensive crisis.

Important Safety Information about JORNAY PM (methylphenidate HCl extended-release capsules)

JORNAY PM
(methylphenidate HCl
extended-release
capsules)

WARNINGS AND PRECAUTIONS

JORNAY PM can cause serious adverse reactions and patients should be monitored for the following:

- Risks to Patients with Serious Cardiac Disease: Sudden death has been reported in patients with structural cardiac abnormalities or other serious cardiac disease who were treated with CNS stimulants at the recommended ADHD dosage. Avoid use in patients with known structural cardiac abnormalities, cardiomyopathy, serious cardiac arrhythmia, coronary artery disease, or other serious cardiac disease.
- Increased Blood Pressure and Heart Rate.
- Psychiatric Adverse Reactions: Including exacerbation of behavior disturbance and thought disorder in patients with a pre-existing psychotic disorder, induction of a manic episode in patients with bipolar disorder, and new psychotic or manic symptoms. Prior to initiating treatment, screen patients for risk factors for psychiatric adverse reactions. If such symptoms occur, consider discontinuing JORNAY PM.
- Priapism: Patients should seek immediate medical attention.
- Peripheral Vasculopathy, including Raynaud's Phenomenon: Observe patients for digital changes during treatment.
- Weight Loss and Long-Term Suppression of Growth in Pediatric Patients: Monitor height and weight.
- Increased Intraocular Pressure (IOP) and Glaucoma: Patients at risk for acute angle closure glaucoma should be evaluated by an ophthalmologist. Closely monitor patients with a history of abnormally increased IOP or open angle glaucoma.
- Onset or Exacerbation of Motor and Verbal Tics, and Worsening of Tourette's Syndrome.

ADVERSE REACTIONS

- The most common ($\geq 5\%$ and twice the rate of placebo) adverse reactions with methylphenidate are decreased appetite, insomnia, nausea, vomiting, dyspepsia, abdominal pain, decreased weight, anxiety, dizziness, irritability, affect lability, tachycardia, and increased blood pressure.
- Additional adverse reactions ($\geq 5\%$ and twice the rate of placebo) in JORNAY PM-treated pediatric patients 6 to 12 years are headache, psychomotor hyperactivity, and mood swings.

DRUG INTERACTIONS

- Antihypertensive drugs: Monitor blood pressure data. Adjust dosage of antihypertensive drug as needed.

Important Safety Information about AZSTARYS (serdexmethylphenidate and dexamethylphenidate capsules)

AZSTARYS
(serdexmethylphenidate
and
dexamethylphenidate
capsules)

WARNING: ABUSE, MISUSE, AND ADDICTION

AZSTARYS has a high potential for abuse, misuse, which can lead to the development of a substance use disorder, including addiction. Misuse and abuse of CNS stimulants, including AZSTARYS, can result in overdose and death.

- **Before prescribing, assess each patient's risk for abuse, misuse, and addiction.**
- **Educate patients and families about risks, proper storage, and proper disposal of any unused drug.**
- **Frequently monitor for signs and symptoms of abuse, misuse, and addiction throughout treatment.**

CONTRAINDICATIONS

- Known hypersensitivity to serdexmethylphenidate, methylphenidate, or product components.
- Concurrent treatment with a monoamine oxidase inhibitor (MAOI), or use of an MAOI within the preceding 14 days.

Important Safety Information about AZSTARYS (serdexmethylphenidate and dexamethylphenidate capsules)

AZSTARYS
(serdexmethylphenidate
and
dexamethylphenidate
capsules)

WARNINGS AND PRECAUTIONS

- Risks to patients with serious cardiac disease: Avoid use in patients with known structural cardiac abnormalities, cardiomyopathy, serious cardiac arrhythmias, coronary artery disease, or other serious cardiac disease.
- Increased blood pressure and heart rate: Monitor blood pressure and pulse.
- Psychiatric adverse reactions: Prior to initiating AZSTARYS, screen patients for risk factors for developing a manic episode. If new psychotic or manic symptoms occur, consider discontinuing AZSTARYS.
- Priapism: If abnormally sustained or frequent and painful erections occur, patients should seek immediate medical attention.
- Peripheral vasculopathy, including Raynaud's Phenomenon: Careful observation for digital changes is necessary during AZSTARYS treatment with ADHD stimulants. Further clinical evaluation (e.g., rheumatology referral) may be appropriate for patients who develop signs or symptoms of peripheral vasculopathy (5.6)
- Long-term suppression of growth in pediatric patients: Monitor height and weight at appropriate intervals in pediatric patients.
- Acute angle closure glaucoma: AZSTARYS-treated patients considered at risk for acute angle closure glaucoma (e.g., patients with significant hyperopia) should be evaluated by an ophthalmologist.
- Increased intraocular pressure (IOP) and glaucoma: Prescribe AZSTARYS to patients with open-angle glaucoma or abnormally increased IOP only if the benefit of treatment is considered to outweigh the risk. Closely monitor patients with a history of increased IOP or open angle glaucoma.
- Motor and verbal tics, and worsening of Tourette's syndrome: Before initiating AZSTARYS, assess the family history and clinically evaluate patients for tics or Tourette's syndrome. Regularly monitor patients for the emergence or worsening of tics or Tourette's syndrome.
- Discontinue treatment if clinically appropriate.

ADVERSE REACTIONS

- The most common adverse reactions ($\geq 5\%$ and twice the rate of placebo) include decreased appetite, decreased weight, nausea, abdominal pain, dyspepsia, vomiting, insomnia, anxiety, affect lability, irritability, dizziness, increased blood pressure, and tachycardia.

DRUG INTERACTIONS

- Antihypertensive Drugs: Monitor blood pressure. Adjust dosage of antihypertensive drug as needed.

Important Safety Information about XTAMPZA ER (oxycodone) extended-release capsules

XTAMPZA ER
(Oxycodone) extended-
release capsules

WARNING: SERIOUS AND LIFE-THREATENING RISKS FROM USE OF XTAMPZA ER

Addiction, Abuse, and Misuse

Because the use of XTAMPZA ER exposes patients and other users to the risks of opioid addiction, abuse, and misuse, which can lead to overdose and death, assess each patient's risk prior to prescribing and reassess all patients regularly for the development of these behaviors and conditions.

Life-Threatening Respiratory Depression

Serious, life-threatening, or fatal respiratory depression may occur with use of XTAMPZA ER, especially during initiation or following a dosage increase. To reduce the risk of respiratory depression, proper dosing and titration of XTAMPZA ER are essential.

Accidental Ingestion

Accidental ingestion of even one dose of XTAMPZA ER, especially by children, can result in a fatal overdose of oxycodone.

Risks From Concomitant Use With Benzodiazepines Or Other CNS Depressants

Concomitant use of opioids with benzodiazepines or other central nervous system (CNS) depressants, including alcohol, may result in profound sedation, respiratory depression, coma, and death. Reserve concomitant prescribing of XTAMPZA ER and benzodiazepines or other CNS depressants for use in patients for whom alternative treatment options are inadequate.

Neonatal Opioid Withdrawal Syndrome (NOWS)

If opioid use is required for an extended period of time in a pregnant woman, advise the patient of the risk of NOWS, which may be life-threatening if not recognized and treated. Ensure that management by neonatology experts will be available at delivery.

Important Safety Information about XTAMPZA ER (oxycodone) extended-release capsules

XTAMPZA ER
(Oxycodone) extended-
release capsules

Opioid Analgesic Risk Evaluation and Mitigation Strategy (REMS)

Healthcare providers are strongly encouraged to complete a REMS-compliant education program and to counsel patients and caregivers on serious risks, safe use, and the importance of reading the Medication Guide with each prescription.

Cytochrome P450 3A4 Interaction

The concomitant use of XTAMPZA ER with all cytochrome P450 3A4 inhibitors may result in an increase in oxycodone plasma concentrations, which could increase or prolong adverse drug effects and may cause potentially fatal respiratory depression. In addition, discontinuation of a concomitantly used cytochrome P450 3A4 inducer may result in an increase in oxycodone plasma concentration. Regularly evaluate patients receiving XTAMPZA ER and any CYP3A4 inhibitor or inducer.

Important Safety Information about BELBUCA (buprenorphine buccal film)

BELBUCA
(buprenorphine buccal
film)

WARNING: SERIOUS AND LIFE-THREATENING RISKS FROM USE OF BELBUCA

Addiction, Abuse, and Misuse

BELBUCA exposes patients and other users to the risks of opioid addiction, abuse, and misuse, which can lead to overdose and death, assess each patient's risk prior to prescribing and reassess all patients regularly for the development of these behaviors and conditions.

Life-Threatening Respiratory Depression

Serious, life-threatening, or fatal respiratory depression may occur with use of BELBUCA, especially during initiation or following a dosage increase. To reduce the risk of respiratory depression, proper dosing and titration of BELBUCA are essential. Misuse or abuse of BELBUCA by chewing, swallowing, snorting, or injecting buprenorphine extracted from the buccal film will result in the uncontrolled delivery of buprenorphine and poses a significant risk of overdose and death.

Accidental Exposure

Accidental exposure of even one dose of BELBUCA, especially in children, can result in a fatal overdose of buprenorphine.

Risks from Concomitant Use with Benzodiazepines Or Other CNS Depressants

Concomitant use of opioids with benzodiazepines or other central nervous system (CNS) depressants, including alcohol, may result in profound sedation, respiratory depression, coma, and death. Reserve concomitant prescribing of BELBUCA and benzodiazepines or other CNS depressants for use in patients for whom alternative treatment options are inadequate.

Neonatal Opioid Withdrawal Syndrome (NOWS)

If opioid use is required for an extended period of time in a pregnant woman, advise the patient of the risk of NOWS, which may be life-threatening if not recognized and treated. Ensure that management by neonatology experts will be available at delivery.

Opioid Analgesic Risk Evaluation and Mitigation Strategy (REMS)

Healthcare providers are strongly encouraged to complete a REMS-compliant education program and to counsel patients and caregivers on serious risks, safe use, and the importance of reading the Medication Guide with each prescription.

Important Safety Information about NUCYNTA ER (tapentadol) extended-release tablets

NUCYNTA ER
(tapentadol) extended-
release tablets

WARNING: SERIOUS AND LIFE-THREATENING RISKS FROM USE OF NUCYNTA ER

Addiction, Abuse, and Misuse

Because the use of NUCYNTA ER exposes patients and other users to the risks of opioid addiction, abuse, and misuse, which can lead to overdose and death, assess each patient's risk prior to prescribing and reassess all patients regularly for the development of these behaviors and conditions.

Life-Threatening Respiratory Depression

Serious, life-threatening, or fatal respiratory depression may occur with use of NUCYNTA ER, especially during initiation or following a dosage increase. To reduce the risk of respiratory depression, proper dosing and titration of NUCYNTA ER are essential. Instruct patients to swallow NUCYNTA ER tablets whole; crushing, chewing, or dissolving NUCYNTA ER tablets can cause rapid release and absorption of a potentially fatal dose of tapentadol.

Accidental Ingestion

Accidental ingestion of even one dose of NUCYNTA ER, especially by children, can result in a fatal overdose of tapentadol.

Interaction with Alcohol

Instruct patients not to consume alcoholic beverages or use prescription or nonprescription products that contain alcohol while taking NUCYNTA ER. The co-ingestion of alcohol with NUCYNTA ER may result in increased plasma tapentadol levels and a potentially fatal overdose of tapentadol.

Risks From Concomitant Use With Benzodiazepines Or Other CNS Depressants

Concomitant use of opioids with benzodiazepines or other central nervous system (CNS) depressants, including alcohol, may result in profound sedation, respiratory depression, coma, and death. Reserve concomitant prescribing of NUCYNTA ER and benzodiazepines or other CNS depressants for use in patients for whom alternative treatment options are inadequate.

Important Safety Information about NUCYNTA ER (tapentadol) extended-release tablets

NUCYNTA ER
(tapentadol) extended-
release tablets

Neonatal Opioid Withdrawal Syndrome

If opioid use is required for an extended period of time in a pregnant woman, advise the patient of the risk of NOWS, which may be life-threatening if not recognized and treated. Ensure that management by neonatology experts will be available at delivery.

Opioid Analgesic Risk Evaluation and Mitigation Strategy (REMS)

Healthcare providers are strongly encouraged to complete a REMS-compliant education program and to counsel patients and caregivers on serious risks, safe use, and the importance of reading the Medication Guide with each prescription.

Important Safety Information about NUCYNTA (Tapentadol) tablets

NUCYNTA
(tapentadol) tablets

WARNING: SERIOUS AND LIFE-THREATENING RISKS FROM USE OF NUCYNTA TABLETS

Addiction, Abuse, and Misuse

Because the use of NUCYNTA tablets exposes patients and other users to the risks of opioid addiction, abuse, and misuse, which can lead to overdose and death, assess each patient's risk prior to prescribing and reassess all patients regularly for the development of these behaviors and conditions.

Life-Threatening Respiratory Depression

Serious, life-threatening, or fatal respiratory depression may occur with use of NUCYNTA tablets, especially during initiation or following a dosage increase. To reduce the risk of respiratory depression, proper dosing and titration of NUCYNTA tablets are essential.

Accidental Ingestion

Accidental ingestion of even one dose of NUCYNTA tablets, especially by children, can result in a fatal overdose of tapentadol.

Risks From Concomitant Use With Benzodiazepines Or Other CNS Depressants

Concomitant use of opioids with benzodiazepines or other central nervous system (CNS) depressants, including alcohol, may result in profound sedation, respiratory depression, coma, and death. Reserve concomitant prescribing of NUCYNTA tablets and benzodiazepines or other CNS depressants for use in patients for whom alternative treatment options are inadequate.

Neonatal Opioid Withdrawal Syndrome (NOWS)

If opioid use is required for an extended period of time in a pregnant woman, advise the patient of the risk of NOWS, which may be life-threatening if not recognized and treated. Ensure that management by neonatology experts will be available at delivery.

Opioid Analgesic Risk Evaluation and Mitigation Strategy (REMS)

Healthcare providers are strongly encouraged to complete a REMS-compliant education program and to counsel patients and caregivers on serious risks, safe use, and the importance of reading the Medication Guide with each prescription.

Important Safety Information about SYMPROIC (naldemedine) tablets

SYMPROIC
(naldemedine) tablets

SYMPROIC may cause serious side effects, including:

- Tear in your stomach or intestinal wall (perforation). Stomach pain that is severe can be a sign of a serious medical condition. If you get stomach pain that does not go away, stop taking SYMPROIC and get emergency medical help right away
- Opioid withdrawal. You may have symptoms of opioid withdrawal during treatment with SYMPROIC including sweating, chills, tearing, warm or hot feeling to your face (flush), sneezing, fever, feeling cold, abdominal pain, diarrhea, nausea, and vomiting. Tell your healthcare provider if you have any of these symptoms

Do not take SYMPROIC if you:

- Have a bowel blockage (intestinal obstruction) or have a history of bowel blockage
- Are allergic to SYMPROIC or any of the ingredients in SYMPROIC. See the Medication Guide for a complete list of ingredients in SYMPROIC. Tell your healthcare provider or pharmacist before you start or stop any medicines during treatment with SYMPROIC

Before you take SYMPROIC, tell your healthcare provider about all of your medical conditions, including if you:

- Have any stomach or bowel (intestines) problems, including stomach ulcer, Crohn's disease, diverticulitis, cancer of the stomach or bowel, or Ogilvie's syndrome
- Have liver problems
- Are pregnant or plan to become pregnant. Taking SYMPROIC during pregnancy may cause opioid withdrawal symptoms in your unborn baby. Tell your healthcare provider right away if you become pregnant during treatment with SYMPROIC
- Are breastfeeding or plan to breastfeed. It is not known if SYMPROIC passes into your breast milk. You should not breastfeed during treatment with SYMPROIC and for 3 days after your last dose. Taking SYMPROIC while you are breastfeeding may cause opioid withdrawal symptoms in your baby. You and your healthcare provider should decide if you will take SYMPROIC or breastfeed. You should not do both
- The most common side effects of SYMPROIC include stomach (abdomen) pain, diarrhea, nausea and vomiting (gastroenteritis)
- Tell your healthcare provider if you have any side effect that bothers you or that does not go away. These are not all the possible side effects of SYMPROIC. Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088

Important Safety Information about SYMPROIC (naldemedine) tablets

SYMPROIC
(naldemedine) tablets

INDICATIONS AND USAGE

SYMPROIC is indicated for the treatment of opioid-induced constipation (OIC) in adult patients with chronic non-cancer pain, including patients with chronic pain related to prior cancer or its treatment who do not require frequent (e.g., weekly) opioid dosage escalation.



CONTRAINDICATIONS

SYMPROIC is contraindicated in:

- Patients with known or suspected gastrointestinal obstruction and patients at increased risk of recurrent obstruction, due to the potential for gastrointestinal perforation
- Patients with a history of a hypersensitivity reaction to Naldemedine. Reactions have included bronchospasm and rash

WARNINGS AND PRECAUTIONS

Gastrointestinal Perforation: Cases of gastrointestinal perforation have been reported with use of another peripherally acting opioid antagonist in patients with conditions that may be associated with localized or diffuse reduction of structural integrity in the wall of the gastrointestinal tract (e.g., peptic ulcer disease, Ogilvie's syndrome, diverticular disease, infiltrative gastrointestinal tract malignancies, or peritoneal metastases). Take into account the overall risk-benefit profile when using SYMPROIC in patients with these conditions or other conditions which might result in impaired integrity of the gastrointestinal tract wall (e.g., Crohn's disease). Monitor for the development of severe, persistent, or worsening abdominal pain; discontinue SYMPROIC in patients who develop this symptom.

Opioid Withdrawal: Clusters of symptoms consistent with opioid withdrawal, including hyperhidrosis, chills, increased lacrimation, hot flush/flushing, pyrexia, sneezing, feeling cold, abdominal pain, diarrhea, nausea, and vomiting have occurred in patients treated with SYMPROIC. Patients having disruptions to the blood-brain barrier may be at increased risk for opioid withdrawal or reduced analgesia. Take into account the overall risk-benefit profile and monitor for symptoms of opioid withdrawal when using SYMPROIC in such patients.



ADVERSE REACTIONS

- The most common adverse reactions with SYMPROIC compared to placebo in two pooled 12-week studies were: abdominal pain (8% vs 2%), diarrhea (7% vs 2%), nausea (4% vs 2%), and gastroenteritis (2% vs 1%).
- The incidence of adverse reactions of opioid withdrawal in two pooled 12-week studies was 1% (8/542) for SYMPROIC and 1% (3/546) for placebo. In a 52-week study, the incidence was 3% (20/621) for SYMPROIC and 1% (9/619) for placebo.

OVERDOSAGE

Single doses of Naldemedine up to 100 mg (500 times the recommended dose) and multiple doses of up to 30 mg (150 times the recommended dose) for 10 days have been administered to healthy subjects in clinical studies. Dose-dependent increases in gastrointestinal-related adverse reactions, including abdominal pain, diarrhea, and nausea, were observed. Single doses of Naldemedine up to 3 mg (15 times the recommended dose) and multiple doses of 0.4 mg (twice the recommended dose) for 28 days have been administered to patients with OIC in clinical studies. Dose dependent increases in gastrointestinal-related adverse reactions, including abdominal pain, diarrhea, nausea, and vomiting, were observed. Also, chills, hyperhidrosis, and dizziness were reported more frequently at 1 and 3 mg doses and hyperhidrosis at the 0.4 mg dose. No antidote for Naldemedine is known. Hemodialysis is not an effective means to remove Naldemedine from the blood.

Important Safety Information about SYMPROIC (naldemedine) tablets

SYMPROIC
(naldemedine) tablets

USE IN SPECIFIC POPULATIONS

Pregnancy:



There are no available data with Naldemedine in pregnant women to inform a drug-associated risk of major birth defects and miscarriage. There is a potential for opioid withdrawal in a fetus when SYMPROIC is used in pregnant women. SYMPROIC should be used during pregnancy only if the potential benefit justifies the potential risk.

Fetal/Neonatal Adverse Reactions

Naldemedine crosses the placenta and may precipitate opioid withdrawal in a fetus due to the immature fetal blood-brain barrier.

Lactation

There is no information regarding the presence of Naldemedine in human milk, the effects on the breastfed infant, or the effects on milk production. Because of the potential for serious adverse reactions, including opioid withdrawal in breastfed infants, a decision should be made to discontinue breastfeeding or discontinue the drug, taking into account the importance of the drug to the mother. If drug is discontinued in order to minimize drug exposure to a breastfed infant, advise women that breastfeeding may be resumed 3 days after the final dose of SYMPROIC.

Pediatric Use

The safety and effectiveness of SYMPROIC have not been established in pediatric patients.

Geriatric Use

Of the 1163 patients exposed to SYMPROIC in clinical studies, 183 (16%) were 65 years of age and over, while 37 (3%) were 75 years and over. No overall differences in safety or effectiveness between these and younger patients were observed, but greater sensitivity of some older individuals cannot be ruled out. In a population pharmacokinetic analysis, no age-related alterations in the pharmacokinetics of Naldemedine were observed.

Hepatic Impairment

The effect of severe hepatic impairment (Child-Pugh Class C) on the pharmacokinetics of Naldemedine has not been evaluated. Avoid use of SYMPROIC in patients with severe hepatic impairment. No dose adjustment of SYMPROIC is required in patients with mild or moderate hepatic impairment.

Non-GAAP Reconciliations

Reconciliation of GAAP Net Income to Adjusted EBITDA

(in thousands, unaudited)

	Three Months Ended June 30,		Six Months Ended June 30,	
	2026	2025	2026	2025
GAAP net (loss) income	\$ (15,051)	\$ 11,983	\$ (555)	\$ 14,400
Adjustments:				
Interest expense	19,519	20,463	35,381	41,253
Interest income	(2,289)	(2,383)	(5,995)	(4,608)
Provision for income taxes	1,519	5,042	5,763	5,747
Depreciation	1,812	1,135	2,275	2,226
Amortization	62,953	55,473	118,426	110,946
Stock-based compensation	14,484	10,818	25,364	22,342
Recognition of step-up basis in inventory	5,417	1,954	5,417	5,431
Executive transition expense	1,393	—	1,393	1,397
Acquisition-related expenses	24,086	935	30,261	2,224
Gain on fair value remeasurement of contingent consideration	—	(358)	—	(1,144)
Total adjustments	\$ 128,894	\$ 93,079	\$ 218,285	\$ 185,814
Adjusted EBITDA	\$ 113,843	\$ 105,062	\$ 217,730	\$ 200,214

Reconciliation of GAAP Operating Expenses to Adjusted Operating Expenses

(in thousands, unaudited)

	Three Months Ended June 30,		Six Months Ended June 30,	
	2026	2025	2026	2025
GAAP operating expenses	\$ 106,594	\$ 73,279	\$ 192,944	\$ 148,916
Adjustments:				
Stock-based compensation	14,484	10,818	25,364	22,342
Executive transition expense	1,393	—	1,393	1,397
Acquisition-related expenses	24,086	935	30,261	2,224
Gain on fair value remeasurement of contingent consideration	—	(358)	—	(1,144)
Total adjustments	\$ 39,963	\$ 11,395	\$ 57,018	\$ 24,819
Adjusted operating expenses	<u>\$ 66,631</u>	<u>\$ 61,884</u>	<u>\$ 135,926</u>	<u>\$ 124,097</u>

Reconciliation of GAAP Net Income to Adjusted Net Income and Adjusted Earnings Per Share

(in thousands, except share and per share amounts, unaudited)

	Three Months Ended June 30,		Six Months Ended June 30,	
	2026	2025	2026	2025
GAAP net (loss) income	\$ (15,051)	\$ 11,983	\$ (555)	\$ 14,400
Adjustments:				
Non-cash interest expense	1,082	1,355	1,901	2,722
Amortization	62,953	55,473	118,426	110,946
Stock-based compensation	14,484	10,818	25,364	22,342
Recognition of step-up basis in inventory	5,417	1,954	5,417	5,431
Executive transition expense	1,393	—	1,393	1,397
Acquisition-related expenses	24,086	935	30,261	2,224
Gain on fair value remeasurement of contingent	—	(358)	—	(1,144)
Income tax effect of above adjustments (1)	(18,959)	(17,871)	(37,588)	(36,608)
Total adjustments	\$ 90,456	\$ 52,306	\$ 145,174	\$ 107,310
Non-GAAP adjusted net income	\$ 75,405	\$ 64,289	\$ 144,619	\$ 121,710
Adjusted weighted-average shares — diluted (2)	39,935,892	39,075,703	40,076,322	39,283,297
Adjusted earnings per share (2)	\$ 1.92	\$ 1.68	\$ 3.67	\$ 3.16

(1) The income tax effect of the adjustments was calculated by applying our blended federal and state statutory rate to the items that have a tax effect. The blended federal and state statutory rate for the three months ended June 30, 2026 and 2025 were 22.9% and 25.7%, respectively; and the blended federal and state statutory rate for the six months ended June 30, 2026 and 2025 were 23.6% and 25.8%, respectively. As such, the non-GAAP effective tax rates for the three months ended June 30, 2026 and 2025 were 17.3% and 25.5%, respectively; and the non-GAAP effective tax rates for the six months ended June 30, 2026 and 2025 were 20.6% and 25.4%, respectively.

(2) Adjusted weighted-average shares - diluted were calculated using the "if-converted" method for our convertible notes in accordance with ASC 260, Earnings per Share. As such, adjusted weighted-average shares – diluted includes shares related to the assumed conversion of our convertible notes and the associated cash interest expense is added-back to non-GAAP adjusted net income. For the three and six months ended June 30, 2026 and 2025, adjusted weighted-average shares – diluted includes 6,606,305 shares attributable to our convertible notes. In addition, adjusted earnings per share includes other potentially dilutive securities to the extent that they are not antidilutive.